Best Reviewers Award for 2019







THE BEST REVIEWER Award has been established to honor our reviewers who made substantial contributions in support of Digestive Endoscopy. Last year, a total of 425 reviewers kindly reviewed manuscripts submitted to the journal, and the Editorial Board wishes to recognize 20 of them for their work. We are bestowing this award on these individuals based on the following criteria:

- 1. Invitation acceptance rate: 80% and over.
- 2. Number of completed reviews (except editorials): five reviews or more; The way to calculate the number of the articles
 - a. Review/Original Articles/How I do it ----- ×1
 - b. Letters/Video Articles/Case Reports ----- ×0.5
 - c. Editorial ----- ×0
- 3. Top 20 reviewers in average scores for review: eight points maximum and two points Minimum, Quality (5-point scale) + Timeliness (3-point scale).

Best Reviewers for 2019

Hiroyuki Aihara

Harvard Medical School, Division of Gastroenterology, Hepatology and Endoscopy, Brigham and Women's Hospital, Boston, MA, USA




Waku Hatta Division of Gastroenterology, Tohoku University Graduate School of Medicine, Miyagi, Japan




Osamu Dohi

Department of Molecular Gastroenterology and Hepatology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto, Japan



Takuto Hikichi Department of Endoscopy, Fukushima Medical University Hospital, Fukushima, Japan



Mitsuru Esaki

Division of Gastroenterology and Hepatology, Department of Medicine, Nihon University School of Medicine, Tokyo, Japan



Keisuke Iwata Department of Gastroenterology, Gifu Prefectural General Medical Center, Gifu, Japan



Yoshihide Kanno

Department of Gastroenterology, Sendai City Medical Center, Miyagi, Japan




Shinichi Nakamura

Department of Gastroenterology, Tokyo Women's Medical University, Tokyo, Japan



Hiroki Sato

Division of Gastroenterology, Graduate School of Medical and Dental Sciences, Niigata University, Niigata, Japan



Koichiro Kawaguchi

Division of Gastroenterology and Nephrology, Faculty of Medicine, Tottori University, Tottori, Japan



Kouichi Nonaka

Department of Digestive Endoscopy, Tokyo Women's Medical University, Tokyo, Japan



Mitsushige Sugimoto

Department of Gastroenterological Endoscopy, Tokyo Medical University Hospital, Tokyo, Japan



Yasuaki Nagami

Department of Gastroenterology, Osaka City University Graduate School of Medicine, Osaka, Japan



Takumi Onoyama

Division of Gastroenterology and Nephrology, Faculty of Medicine, Tottori University, Tottori, Japan




Sho Suzuki

Division of Gastroenterology and Hepatology, Department of Medicine, Nihon University School of Medicine, Tokyo, Japan



Mamoru Takenaka

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osaka, Japan



Tomonori YanoDepartment of Gastroenterology and Endoscopy, National Cancer Center Hospital



Yosuke Tsuji

Department of Gastroenterology, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan



Masao Yoshida Division of Endoscopy, Shizuoka Cancer Center, Shizuoka, Japan



Yasushi Yamasaki

Department of Gastroenterology, Okayama University Hospital, Okayama, Japan

